



SiiLK MesoFAT

V-LINEFACE
S-LINE SHAPE

SiiLK MesoFAT VL

MesoFAT VL is an effective solution in fat reduction, using *Deoxycholic Acid* which has been proven its safety and efficacy by pharmaceutical clinical evaluation.



MAIN INGREDIENTS

DEOXYCHOLIC ACID

Deoxycholic Acid is widely used as an alternative to liposuction, to reduce subcutaneous fat. The minimal invasive technique Lipolysis Injection using Deoxycholic Acid is safe and approved by the Food and Drug Administration (FDA).

Deoxycholic Acid naturally found in human body as a component of bile acid, that has a vital role in emulsification & digestion of fat in intestine.

Deoxycholic Acid disrupts fat cells membrane physically, causes the death of adipocytes, and the size reduction. Meanwhile, inflammatory cascade resulting from the necrosis cells, causing fibroblasts stimulation, and produces more collagen (neo-collagenesis). The collagen production improves tissue quality, firm skin texture, and renew the skin.

PEPTIDES

After the subcutaneous layer lost many of fat cells, it leaves empty space and causes sagging appearance. Peptides take many important roles in our body, including the skin. It will promotes skin tightening, skin conditioning, improves lines & wrinkles, and rejuvenation, thus helps the skin after lipolysis procedure to regain its elasticity and prevent the sagging appearance.

OLD LIPOLYSIS

MODE of ACTION	Lipolysis.
MAIN RESULT	Not destructing fat cells. Only short-term temporary effect by reducing fat size.
AFTER USE	Rebound phenomenon (as the fat cells are in famine condition).

SIIILK MESOFAT

MODE of ACTION	Dual Action (Lipolysis & Adipocytolysis)
MAIN RESULT	Fat cells disrupted & destroyed. Long-term effects.
AFTER USE	No rebound due to disruption of fat membrane, total fat cells reduction, and neo-collagenesis to repair the skin.

SIILK MESOFAT VL

CHEEK

3 - 10 points per side (1.5cm - 2.5cm distance)
0.1 ml - 0.2ml per point injection
Injection depth (subcutaneous layer : 4-8mm)

DOUBLE CHIN


3 - 8 points (2cm - 4cm distance)
0.1 ml - 0.4ml per point injection
Injection depth (subcutaneous layer : 6-10mm)

JAWLINE AREA


5 - 8 points per side (1.5cm - 2cm distance)
0.1 ml - 0.15ml per point injection
Injection depth (subcutaneous layer : 2-6mm)

SUGGESTION

Minimum 3-4 sessions in 10-14 days interval



INDICATED AREA & PROTOCOL



SIILK MESOFAT SL

ARMS, BACK AREA, WAIST

2cm - 2.5cm distance
0.2ml - 0.3ml per point injection
Injection depth (subcutaneous layer : 6-10mm)

ABDOMEN, THIGHS, BUTTOCKS

2cm - 3.5cm distance
0.2ml - 0.4ml per point injection
Injection depth (subcutaneous layer : 8-12mm)

BUTTOCKS UNDER LINE

1.5cm - 2cm distance
0.1 ml - 0.2ml per point injection
Injection depth (subcutaneous layer : 6-10mm)

SUGGESTION


Minimum 5-6 sessions in 10-14 days interval



SiiLK MesoFAT SL

MesoFAT SL is an effective and safe fat dissolving system with **Dual Action** of Deoxycholic Acid (Adipocytolysis) and Soybean Extract (Strong Lipolysis) combination.

MODE of ACTIONS



ADIPOCYTOLYSIS


Deoxycholic Acid disrupts fat cells membrane, causing fat cells necrosis, reduce fat cells & fat size.

LIPOLYSIS

Deoxycholic Acid combined with PhosPhatidylCholine (PPC) extracted from soybean emulsifies and destroys the fat, then removes it into blood stream to be naturally discharged through metabolism.

NEO-COLLAGENESIS

The death necrotic fat cells activates inflammatory cascade, causing fibroblast stimulation to promote neo-collagenesis. Empty space left by fat cells destruction will be improved and repair by new healthy collagen.



- **FAT REDUCTION**
- **NO REBOUND**
- **LONG TERM**
- **SKIN TIGHTENING**

*'Slimming Solution,
S-Line Body Shape'*

----- MesoFAT SL



COMPOSITION

Deoxycholic Acid
Soybean Extract
Peptides

PACKAGE

8ml x 5vials

STORAGE

Store in a cool room temperature, avoid sunlight.
After opened, put inside refrigerator under 10-15°C
and use it within 15 days.

CAUTION

Shake well before use, subcutaneous procedure only.
Do not use to patients with history of allergic to
soybeans, beans, and all the derivations.
Do not use to pregnant patients or during lactation.
In the presence of side effects, stop usage and
contact your dermatologist immediately.



SiiLK MesoFAT

COMPOSITION

Deoxycholic Acid
Peptides

PACKAGE

8ml x 5vials

STORAGE

Store in a cool room temperature, avoid sunlight.
After opened, put inside refrigerator under 10-15°C
and use it within 15 days.

CAUTION

Do not use to pregnant patients or during lactation.
In the presence of side effects, stop usage and
contact your dermatologist immediately.





 +82-(0)10-6365-3328

 info@beaumed.co.kr

 www.beaumed.co.kr



#BeautyMadeby BEAUMED

